

FACT SHEET ON BISPHENOL A


Cancer Association of South Africa (CANSA)

August 2010


Compiled by Dr C Albrecht – Head of Research

WHAT IS BISPHENOL A? ^{1,2,3,4,5}

- Bisphenol A (abbreviated BPA) is a high-production volume, artificial, industrial, man-made molecule that was first synthesised in 1891 by A.P. Dianin.¹ It is an ever-present environmental contaminant with established endocrine disruptor properties.¹⁰
- It is a relatively small symmetric, organic compound (see below) with a molecular weight of 228. It is a white powder and surprisingly has been found to mimic the female hormone, oestrogen, by binding to the oestrogen receptor. It is one of the best examples of a so-called endocrine-disrupting compound (EDC) because it binds to oestrogen-receptors in the cell in a haphazard, uncoordinated way - disrupting the highly ordered working of the normal hormone, oestrogen, especially during pregnancy, and causing various health defects, i.e. it is a pseudo-oestrogen.^{1,6}
- BPA was investigated in the 1930's as a possible synthetic oestrogen to be used for miscarriages. However, it was not developed further because another synthesized compound, diethylstilbestrol (DES), was found in 1940, which turned out to be a much more powerful oestrogen substitute in humans than BPA. Contrary to expectations DES was later taken off the market in 1970 when it was found to be linked to vaginal adenocarcinoma (cancer) and foetal malformations in daughters of 7 million mothers who used the drug while pregnant. This was a pharmaceutical disaster and in retrospect was an early warning signal for similar toxic properties confirmed for BPA 30 – 40 years later.⁷
- In 1953 Daniel Fox and Hermann Schnell synthesized polycarbonate for the first time using BPA and phosgen⁸. Bisphenol A is mainly used as a bifunctional monomer in the manufacture of polycarbonate plastic and epoxy resins and as an antioxidant in PVC.



BPA as a single molecule




BPA as part of a polycarbonate polymer⁹


WHY IS BPA A MATTER OF CONCERN?

- Exposure to BPA has been linked to increased breast cancer risk, early puberty, obesity, infertility in males and females, brain dysfunction, thyroid dysfunction, heart disease, diabetes and prostate cancer⁹⁻¹⁹
- Evidence has been found that BPA acts epigenetically, at low concentrations, to permanently silence critical genes during development in the womb with as yet unknown consequences^{12,13}
- Substantial evidence indicates that exposure to BPA during early development may increase breast cancer risk. This is supported by studies showing that BPA activates genes involved in growth in mammary cells^{12,14}
- Children may be particularly vulnerable to BPA²
- BPA was detected in the urine of 92% of the U.S. population in 2003-2004 and it is most likely to be in an average South African's urine as well²⁰
- BPA is not essential and provides no known benefit to human health and appears to be harmful to human health
- In 2007 Prof Fred vom Saal of the University of Missouri-Columbia (World expert on BPA) found that very low level exposure of BPA (1 nM) harms the prostate²²
- Between 1997 and 2008, over 100 publications linked very low level exposure of BPA with prostate damage, breast and prostate cells predisposed to cancer, decline in testosterone, changes in breast tissue that predispose cells to hormones and carcinogens, early puberty, behavioural problems and other effects^{22,26}
- These effects have been found at BPA concentrations up to 25 times lower than the U.S. Environmental Protection Agency's (EPA) "safe" dose of 50 micrograms per kilogram (body weight) per day^{22,23}
- 38 Independent BPA scientists (Chapel Hill Panel) completed an assessment of BPA safety in 2007 and concluded that BPA exposure at current levels presents a clear risk to human health²⁴
- In a New York Times editorial (May 20th 2008), a ban on BPA in the U.S. is called for in terms of items such as baby bottles and cups²⁵
- BPA induces a profile of tumour aggressiveness in high-risk cells from breast cancer patients¹²
- Prenatal exposure to BPA induces early cancerous changes in the breast tissue of rats¹⁰⁻¹⁴

EFFECT OF LOW DOSE BPA ON BREAST TISSUE.³



Control milk ducts in mouse breast tissue



Mice treated with 25 ng BPA/kg body weight per day.

Milk duct growth dramatically stimulated


HOW HAS THE WORLD REACTED?

- **Canada.** On April 18, 2008, the Canadian Government **moved to ban polycarbonate infant bottles** as it officially declared one of its ingredients (bisphenol A) toxic²¹
- **Denmark.** From the 1st July 2010 all food contact materials for young children (ages 0-3 yrs), including baby bottles made of polycarbonate, will be banned in Denmark. This was due to concerns of food safety experts that BPA could inhibit learning capacity of children⁴¹
- **Belgium.** March 2010. Legislation to ban BPA in food contact plastics proposed
- **New York.** August 3, 2010. Sales of baby bottles, sippy cups, pacifiers and straws that contain plastic made from Bisphenol A will be banned in the state under law. Also banned in Minnesota, Chicago and Connecticut
- **United States:** April 2010. Annual report of President Obama's President's Cancer Panel states "...the available evidence argues for a precautionary approach to BPA". March 2010. The Environmental Protection Agency (EPA) declared BPA a chemical of concern". The FDA promises to respond by end November 2010 and so does European Food safety Authority (EFSA)

- **France:** 10 August 2010. 9 Senators call for BPA ban in food containers such as baby bottles⁴⁰

WHERE IS BISPHENOL A FOUND?

- Hard, clear, near shatterproof plastic – polycarbonate.
- Epoxy resin coating in metal food cans^{30, 35}
- Plastic bottles and cups for babies. Heating these bottles releases up to 55-times more BPA^{28,29}
- Precursor of flame-retardant – tetrabromobisphenol A
- Bicycle helmets
- Car safety seats
- Water coolers
- Medical devices
- CD's, credit cards, cell phones, computers, cars
- Sports equipment
- Household electronics
- Some drinking water^{30, 33, 35,}
- Global production in 2003 was 2.2 million metric tones in 1999 (This is 2 200 000 000 kg or 328 grams for every man, women and child on earth per year!)¹
- Recently BPA has been reported on cash receipts printed with the heat process such as those dispensed by ATM's. In some cases receipts contained 250 to 1000 times more BPA than is found in baby bottles³⁸
- There is ever-present human exposure to BPA which is a pollutant across the whole planet only dropping to low levels over the poles^{42,36}





BPA on receipts

WHAT SOURCE OF BPA POSES THE GREATEST THREAT?

- Exposure to boiling water increased the rate of BPA migration by up to 55-fold from polycarbonate drinking bottles, as especially those used by babies^{28,29}
- BPA found in the liquid contents of various tinned food products^{30,35}
- BPA rubbing of certain receipts
- Small amounts in the air, water and soil across the planet

HOW CAN THE POTENTIAL CANCER RISK OF BISPHEENOL A BE REDUCED?

CANSA advocates:

- Banning the production of BPA-containing products from which BPA can migrate into food and drink in concentrations known to cause adverse biological effects especially in children – 0-3 years old
- The developing of BPA-free alternatives for baby drinking bottles, the lining of tin cans and outdoor water bottles
- Finding an alternative for receipt paper containing BPA

- Ensuring that drinking water is BPA free
- The investigation of multiple uses of BPA and ascertaining which uses may pose a health threat to man
- Lobbying the government to protect citizens from untoward exposure to BPA
- Not putting polycarbonate plastic food containers in the microwave
- Avoiding plastic utensils with a number 7 in a triangle and the letters PC underneath the triangle. This indicates polycarbonate

LACK OF CONSENSUS:

- There is no consensus in the world about the potential dangers of low dose BPA. The American Chemistry Council believes that BPA is a good product and safe.³⁷ The FDA has expressed 'some concern' but advocates no changes in the status quo at present. **Nevertheless, the public and the baby bottle industry are turning away from BPA in order to be safe, rather than sorry**

SELECTED QUOTATIONS:

“We know a women’s lifetime risk of breast cancer is directly linked to her lifetime exposure to estrogen – both natural and synthetic estrogen. It’s outrageous that manufacturers of some baby bottles are exposing little girls to BPA, a synthetic plasticizer that mimics estrogen, and possibly increasing that little girl’s risk of breast cancer later in life, especially when safe alternatives are available.”

Janet Nudelman - Director, Program and Policy for the Breast Cancer Fund

BPA alone is “worth at least a million dollars every hour”.

Wade Welshons - University of Missouri

“If a chemical is biologically active and interacts with our receptors, it’s probably no good. Ban it.”

Sheldon Krinsky - Tufts University

“If I were a pregnant woman, I would try hard to avoid exposure to BPA”

Randy Jirtle

REFERENCES and SELECTED READING ON BPA:

1. **Bisphenol A** from Wikipedia. http://en.wikipedia.org/wiki/Bisphenol_A (Comprehensive and up to date.)
2. **“Body Toxic”**. Nena Baker, North Point Press, New York, 2009. (How the hazardous chemistry of everyday things threatens our health and well-being).
3. **“Slow Death by Rubber Duck”**, Rick Smith and Bruce Lourie, Alfred A. Knopf Canada, 2009. (How toxic chemistry of everyday life affects our health)
4. **“The Secret History of the War on Cancer”**, Devra Davis, Basic Books, 2009. (A wake-up call for all those who have accepted the poisons of our age of plenty without a blink.)
5. **“The Hundred-Year Lie”**, Randall Fitzgerald, A Plume Book, 2007. (How to protect yourself from the chemicals that are destroying your health.)
6. **Estradiol and Bisphenol A stimulate androgen receptor and oestrogen receptor gene expression in fetal mouse prostate mesenchyme cells**, Richter CA, et al., Environ health Perspect., 2007, **115**, 902-908.
7. **Pharmaceutical Teratogens, diethylstilbesterol**, <http://teratology.org/jfs/Pharmaceutical.html>
8. **History of the invention of polycarbonates**, <http://novattro.net/polycarbonate/history/>
9. **Bisphenol A induces a profile of tumour aggressiveness in high-risk cells from breast cancer patients**, Cancer Res., 2008, **68**, 2076 – 2080.
10. **Proteomic analysis in mammary glands of rat offspring exposed in utero to bisphenol A**, J Proteomics., 2010, **73**, 1241 – 1253.
11. **Prenatal bisphenol A exposure induces preneoplastic lesions in the mammary gland in Wistar rats**, Durando M, et al., Environ Health Perspect., 2007, **115**, 80 – 86.
12. **Epigenetic influences of low-dose bisphenol A in primary human breast epithelial cells**, Weng YI et al, Toxicol Appl Pharmacol., 2010, Jul 30 Epub (Ahead of print).
13. **Epigenomic disruption: The effects of early development exposures**, Birth Defects Res A Clin Mol Teratol, 2010, Jun 21 [Epub ahead of print).

14. **In utero exposure to Bisphenol A shifts the window of susceptibility for mammary carcinogenesis in the rat.**, Betancourt AM et al., Environ Health Perspect., 2010, Jul 30 [Epub ahead of print].
15. **Perinatal exposure to environmental estrogens and the development of obesity**, Newbold RR, et al., Mol Nutr Food Res, 2007, 51, 912-917
16. **Semen quality and sperm DNA damage in relation to urinary bisphenol A among men from an infertility clinic**, Meeker JD et al. Reprod Toxicol, 2010, [Epub ahead of print].
17. **Bisphenol A interferes with synaptic remodeling**, Hajszan T and Leranthy C., Front neuroendocrinol, 2010, Jul 6 [Epub ahead of print].
18. **Toxic environment and obesity pandemic: Is there a relationship?**, Latini G et al, Italian Journal Pediatrics, 2010, 36, <http://www.ijponline.net/content/36/1/8>
19. **In vitro effects of bisphenol A on developing hypothalamic neurons**, Iwakura T et al., Toxicology, 2010, 272, 52-58.
20. **Exposure of the U.S. population to bisphenol A and 4-tertiary-octylphenol: 2003 – 2004**, Calafat, AM et al., 2008, Environ Health Perspect., 116, 39- 44.
21. **Government of Canada Takes Action on Another Chemical of concern: Bisphenol A**, http://www.hc-sc.gc.ca/ahc-asc/media/nr-cp/2008/2008_59_e.html

Canada bans Plastic Bottles tied to health concerns, New York Times , April 18, 2008.
22. **An extensive new literature concerning low-dose effects of bisphenol A shows the need for a new risk assessment**, Vom Saal, FS and Hughes C, Environ Health Perspect.,2005, 113, 926-933.
23. **Many studies confirm BPA's low-dose toxicity across a diverse range of toxic effects**, Environmental Working Group Report. <http://www.ewg.org/node/20941>
24. **Chapel Hill bisphenol A expert panel consensus statement: Integration of mechanisms, effects in animals and potential to impact human health at current levels of exposure**, Reproductive Toxicology, 2007, 24, 131-138.
25. **Concerns about BPA plastic**, The New York Times Editorial, May 20, 2008.
26. **CERHR Expert Panel Report for Bisphenol A**, National Toxicology Program, U.S. Department of Health and Human Services, 26-11-2007.
27. **Bisphenol A is released from polycarbonate drinking bottles and mimics the neurotoxic actions of estrogen in developing cerebellar neurons**, Hoa HL et al., Toxicol Lett.,2008, 176, 149-156.

28. **Migration of bisphenol A from polycarbonate baby bottles under real use conditions**, Maragou NC et al., Food Addit Contam., 2008, **25**, 373 – 383.
29. **Bisphenol A migration from polycarbonate baby bottle with repeated use**, Nam SH, Seo YM and Kim MG, Chemosphere, 2010, **79**, 949-952,
30. **Xenooestrogens released from lacquer coatings in food cans**, Brotons, JA et al., Environ Health Perspect., 1995, **103**, 608 – 612.
31. **Determination of bisphenol A in drinking water system components by liquid chromatography-tandem mass spectrometry**, Lin S et al., Wei Sheng Yan Jiu, 2008, **37**, 85-87.
32. **Derivation of a bisphenol A oral reference dose (RfD) and drinking-water equivalent concentration**, Willhite CC et al., J Toxicol Environ Health B Crit Rev., 2008, **11**, 69-146.
33. **A national reconnaissance for pharmaceuticals and other organic wastewater contaminants in the United states –II. Untreated drinking water sources**, Focazio MJ et al., Sci Total Environ., 21 April 2008.
34. **Occurrence and temporal variations of the xenooestrogens bisphenol A, 4-tert-octylphenol, and tech, 4-nonylphenol in two German wastewater treatment plants**, Hohne C and Puttmann W, Environ Sci Pollut Res Int., May 28 2008.
35. **Bisphenol A in canned food products from Canadian markets**, J Food Prot, 2010, **73**, 1085 – 1089.
36. **Bisphenol A: Invisible pollution**, Groff T, Curr Opin Pediatr, 2010, **22**, 524-529.
37. **Arguments in favour of BPA by the Americal Chemistry Council** - <http://factsaboutbpa.org/>
38. **High levels of BPA on receipts**, <http://www.foodsafetynews.com/2010/08/according-to-new-laboratory-tests/>
39. **Environmental causes of cancer: endocrine disruptors as carcinogens**, Soto AM and Sonnenschein C, Nat Rev Endocrinol, 2010, **6**, 363-370.
40. **Bisphenol A ban tabled in French parliament**, <http://www.foodqualitynews.com/Legislation/Bisphenol-A-ban-tabled-in-French-parliament>
41. **Denmark bans bisphenol A in food packaging for young children**, <http://www.foodproductiondaily.com/Quality-Safety/Denmark-bans-bisphenol-A-in-food-packaging-for-young-children>
42. **Ubiquity of bisphenol A in atmosphere**, Fu P, Kawamura K, Environ Pollut., 2010, Jul 31 [Epub ahead of print]